

Microbiome agility facing the challenge of modern broiler production

Christophe Bostvironnois, DVM – Chr. Hansen, Hoersholm, Denmark
 Randi Lundberg, DVM, PhD – Chr. Hansen, Hoersholm, Denmark



Introduction

In recent years, knowledge about the gut microbiome has increased tremendously in both humans and animals. Actually, for humans, the microbiome is approximately 1.5 kg of bacteria that we are carrying every day. These bacteria and other microorganism part of the gut microbiota are mainly favorable to humans and are acting at the interface of the external world and our body. There is much research leading to the conclusion that this microbiome has an unbelievable impact, not only on the digestive system, but also on the rest of the human or animal body. It influences the immune system, the central nervous system and can as such even affect mood and behavior.

In poultry, we focus a lot on achieving a healthy gut since a well-functioning gut is the key engine for growth, health and welfare of the birds. Agility of the microbiome is closely linked to gut health. That is why we want to take very good care of the chickens' gut microbiota as part of our management practices.

Microbiome Agility comes from a diverse and functional potential of the microbiota

Microbiome Agility can be defined as the ability of the intestinal microorganisms to adapt to changing conditions in order to maintain microbial balance and productivity. Microbial balance leads to immune balance, which is crucial for the animal to overcome intestinal challenges. Microbial productivity means that the gut bacteria are constantly ensuring optimal nutrient utilization and production of compounds that are vital to the animal.

During the life of the birds, they will experience unavoidable changing conditions, such as diet shifts or exposure to pathogens. This is Why Microbiome Agility is crucial to maintain that important microbial balance and productivity. The former way to manage pathogens and thus the microbiota by using massive amounts of chemicals (zinc, copper, antibiotics) is over. At the same time, it is still a dream to get rid of pathogens in modern poultry production, so viable alternatives are needed. Additionally, the use of highly digestive substances is becoming increasingly inefficient from an economic standpoint. As a result, the strategic nutritional formulation is moving to less security and safety for the birds. We are now seeing more use of wheat, Distiller's Dried Grains with Solubles (DDGS), sunflower meal, rapeseed meal or even peas instead of soy meal and corn. So not only do we have less chemicals accepted by the consumers and authorities, but also economically we are looking for alternative raw materials sourced, for example, as a by-product. Ultimately, the pathogens are still there, and the physiology of

the birds is even more challenged with higher growth rate and higher feed intakes. Genetic enhancements are ongoing, but eventually it takes time for birds to develop and mature their physiological enzyme system, immune system, digestive structure of villi and microvilli.

Graph 1. Microbiome Agility can be defined as the ability of the intestinal microorganisms to adapt to changing conditions in order to maintain microbial balance and productivity.




Maintaining microbiome agile by making sure the digestive tract is populated with a diverse and functional microbiota is crucial for the chicken to withstand the challenges as a production animal. We need to look at solutions which can help to achieve this goal. The former way of nutritional and farm management was to get rid of all pathogens. Nowadays we are looking at more holistic approaches for obtaining a plastic gut microbiota that provides protection against pathogens and shapes the digestive physiology and the immune system in beneficial ways. Agility depends on the diversity of the microbiome, so it is not enough just to have the presence of a few different beneficial microbes, like cultivating hectares of only one crop. Too little so-called alfa diversity will make the microbiota more susceptible to challenges, and thus highly unstable. One can say that if we introduce changes in the life of the chicken (like feed transition as a key example), there is a lack of a plan B to adjust to this new reality because there are simply too few good

bacteria to carry the torch. On the other hand, having a good diversity of different microbes with beneficial properties allows to adjust, adapt and take advantage of the different attributes of the variety of favorable species and strains. A lack of diversity makes the whole digestive system extremely fragile and makes it very susceptible to any change.

In that sense, the ideal is to achieve a microbiota in the gut which looks like a forest: many vegetal, a lot of diversity of herbs, fungi and plants, and ultimately huge trees which are dominating the rest of the vegetal, making it difficult for the bad herbs to become dominant. This is achieved by the natural competition on sun, earth and water and their established dominance (Graph 2).

Graph 2. Maintaining Function and diversity, a paradigm shift in modern broilers




A high diversity provides more functional potential to the microbiota. The more diverse, the more plastic and agile it will be. The favorable bacteria will be so dominant that they will de facto prevent pathogenic bacteria to multiply. Just like the trees in the forest are dominating the other vegetal species, the established dominance dictates the natural competition.

A challenging period in the life of a chicken

When we consider the life of a chicken, not much is currently done to achieve microbiome plasticity and agility and many challenges are rising at the horizon. Actually, quite a lot of different events are happening due to management practices, which can affect the microbiome agility. Let's have a look at these challenges (Graph 3).

Graph 3. Multiple Risks of losing Microbiome Agility during the Life of a chicken



First days: Feed intake and first bacterial implementation challenge

Feed can be considered as the fuel of the digestive engine. When we consider the chick weight and the amount of feed it needs to ingest every day, the challenge is very high. For instance, day 1, the ratio between the feed intake and the weight is superior to 30%! For a 4kg human baby, this would represent the ingestion of 1.2kg of food and something like 2 liters of water! Feed, water, heat and oxygen are the key elements for bacterial establishment during the first days of life. Since the digestive tract of the newly hatched chicken is virtually sterile, or comparable to a desert with respect of vegetation, it is easy to understand that there is a massive opportunity for any pathogenic bacteria to grow in such an environment without competition from good bacteria. Much research is showing that this first implementation is based on chance. Any bacteria present in the environment of the day-old birds is likely to be present in the gut of the chick. It explains why there is typically a gram-negative challenge (*E coli*, *Salmonella*) early in life. These bacteria have the capacity to multiply quickly and to take advantage of the new environment: a desert where feed, water, heat and oxygen are in abundance, but no other vegetation to compete these resources. Besides causing pathogenic infections, these bacteria take up the ecological niches in the digestive tract, making it more difficult for more beneficial bacteria to establish. As a result, it is likely that it will lead to a less plastic and agile microflora which will require antibiotic treatment to keep the balance in check and provoke subsequently mortality.

Second and third week: the microbiota shift

The feed intake challenge is diminishing in the second and third weeks of the chick's life (still 20% of feed intake over the individual weight). However, there are other reasons making these weeks very challenging for the microbiome. A natural part of the microbiome establishment is the lower availability of oxygen in the gut happening after the aerotolerant species have established. At this point in time, the strict anaerobic bacteria can colonize. The strict anaerobes are favorable for the bird and gram-positive. However, one of them is *Clostridium perfringens*, a pathogen causing necrotic enteritis. Typically, these bacteria reside at the very end of the digestive tract, in the caeca where oxygen is very low, but they can take advantage of the retro-peristaltic movements of the chicken gut to move up and start proliferation in the small intestine. *Clostridium perfringens* proliferates, leads to an unhealthy balance of the microbiota called dysbiosis, eventually resulting in necrotic enteritis between day 14 and day 18. This is a very typical observation which is also simultaneous to a brutal feed transition between starter and grower diet. It can be even more dramatic in case of vaccination during this period.

Fourth and fifth week: Coccidia and mucus production

These two weeks are dominated by the *Eimeria* challenges which will occur in all farms, regardless of the vaccination program or chemical/ionophore-based program in place. There will be sporozoites of coccidia (either wild or vaccinal strains) which will be in close contact with the epithelial cells of the intestine. As a matter of defense, the first line of protection will be an increased production of mucus by the goblet cells. The aim will be to build and reinforce a wall of mucus to protect the intestine from the parasites' attack. Unfortunately, some pathogens may take advantage of this mucus production. We call them mucolytic bacteria. Not only are they able to survive and multiply within the newly produced layers of mucus but also, they can use the mucus as a source of energy for their own development. As a result, they multiply in situ, in very close contact with the top of the epithelial cells. It is obvious that it will lead to a digestive syndrome so-called dysbacteriosis or dysbiosis. Ultimately, this may be even more damaging when there is the production of digestive toxins such as alpha or net-B toxins produced by *Clostridium perfringens*.

At the same time, this is a period where the broiler producer wants to make the most profit out of its production. The feed intake and the growth rate are becoming very high. This is the period when feed efficiency matters the most. Thus, it is even more critical to maintain a healthy and agile microbiome. Each day of a poorly balanced microbiota can lead to a potential loss of 80 to 100g of body weight/bird. That is equal to 0.08 to 0.1 €/bird of revenue. Considering production of 10,000,000 birds/year, the loss becomes a significant amount of money (800,000 to 1,000,000 €).

In addition, this is a period of intense feed intake stress for the microbiome: brutal transition from grower to finisher feed followed by another transition from finisher to withdrawal feed. When the market needs are there, the poultry producer may use the thinning method when part of the flock will be slaughtered while the rest will stay in place for further growing time in the barn. This is requiring a brutal stop in feeding and drinking the animals for carcass quality reasons. This leads to a complete disruption of microbiome. In a couple of hours, the intestinal content and thus the nutrients, are reduced to a minimal presence, as well as the water content of the digestive tract. When the thinning is finished, the birds typically turn to an over consumption of feed in a short period of time. The microbiome, already challenged by the fasting period now has more feed to be used for rapid development. How do we manage this? Well, we pay a little attention to this practice from a microbiome standpoint. However, there is no doubt that this is completely disturbing the agility and plasticity of the microbiome.

What do we need to do to achieve and maintain Microbiome Agility?

The modern poultry industry should consider Microbiome Agility as a key goal to achieve sustainable poultry production in the future. This is probably the very early stage of a concept which will revolutionize our approach to the intestinal content of the chicken. Whatever the tools, techniques, approaches we will have in the future, we need to achieve three key goals:

1. To seed the gut with favorable bacteria at the right time, probably as early as possible: day 1 and or even in ovo. Seeding with good probiotic bacteria helps prevent overgrowth of bad bacteria. At the same time, it paves the way to the establishment of a well-balanced, well-functioning and highly diverse microbiota. The early seeding and colonization is the first strategic component of the Microbiome Agility.
2. The second is the need to support the multiplication of the favorable bacteria and to allow them to dominate most of the microbiota. This can be achieved by creating a favorable micro-environment in the gut. Well selected sporulated probiotics are dedicated to creating such environment. Other probiotic strains help by competing chemically via bacteriocins or through enzymes that works against potential pathogenic bacteria.
3. Thirdly, there is a need to decrease, as much as possible, undigestible substrates that are potential feed for fermenter and unfavorable bacteria. This ideal substrate for pathogenic bacteria can subsequently lead to enteritis and microbiota unbalance. Recent research in the probiotic world has shown that very specific strains are able to behave as an in-situ enzyme factory, releasing them directly into the gut. Thus, these favorable bacteria can transform the indigestible intestinal content into digestible and valuable nutrients for the birds and not for the microbiota.

Conclusions


We are at the very beginning of a new era for knowledge and understanding of the chicken's microbiome. We are scratching the surface of something bigger. However, what we know from other species, including humans, can make us optimistic. The progresses will be fast. It will undoubtedly allow the development of new solutions better accepted by modern poultry production and by consumers.

References

- Albazaz R.I., 2014, *Microflora of Digestive Tract in Poultry*, <http://dogadergi.ksu.edu.tr/download/article-file/212071>
- Choct M., 2009, *Managing gut health through nutrition*, <https://www.tandfonline.com/doi/pdf/10.1080/00071660802538632?needAccess=true>
- Collier, CT, Van der Klis, JD, Deplancke, B, Anderson, DB & Gaskins, HR 2003, 'Effects of tylosin on bacterial mucolysis, *Clostridium perfringens* colonization, and intestinal barrier function in a chick model of necrotic enteritis' *Antimicrobial Agents and Chemotherapy*, vol 47, no. 10, pp. 3311-3317. DOI: 10.1128/AAC.47.10.3311-3317.2003, <https://www.ncbi.nlm.nih.gov/pubmed/14506046>
- Collier C.T., 2007, *Coccidia-induced mucogenesis promotes the onset of necrotic enteritis by supporting Clostridium perfringens growth*, <https://experts.illinois.edu/en/publications/coccidia-induced-mucogenesis-promotes-the-onset-of-necrotic-enter>
- De Gussem M., 2010, *Coccidiosis in poultry: review on diagnosis, control, prevention and interaction with overall gut health*, <https://www.cabi.org/isc/FullTextPDF/2009/20093257328.pdf>
- Gao et Al, *Feed-additives probiotics accelerate yet antibiotics delay intestinal microbiota maturation in broiler chicken (2017) 5:91.*
- Ginter A., 2007, *Summary of Clostritest® profiles in chicken production*, <http://www.journees-de-la-recherche-cunicole.org/PDF/P10-BOSTVIRON-NOIS%20version%20def.pdf>
- Kers J.G. et Al, *Host and Environmental Factors Affecting the Intestinal Microbiota in chickens*, *Frontiers in Microbiology (2018) 9:235.*
- Klasing K.C., 2007, *Nutrition and the immune system*, <https://www.tandfonline.com/doi/full/10.1080/00071660701671336?scroll=top&needAccess=true>
- Mancabelli L. et Al, *Insights into the biodiversity of the gut microbiota of broiler chickens*, *Environmental Microbiology (2016) 18(12), 4727-4738*
- Van Immerseel, 2011, *Necrotic enteritis in broilers: an updated review on the pathogenesis*, <https://www.ncbi.nlm.nih.gov/pubmed/21812711>



Some words on the author



Christophe Bostvironnois is a Doctor of Veterinary Medicine from National Veterinary School of Lyon (France). He oversees the Poultry portfolio commercial development for Chr Hansen. Prior to that, he held different roles in the feed industry and the animal health industry since he joined the poultry world in 1996. He made since then several publications and conferences on the intestinal and respiratory physiology and diseases of chickens, turkeys and layers. He is also president of the French Branch of the World Poultry Science Association.